arXiv:1301.0406v1 [hep-ex] 3 Jan 2013

Prospects for time-dependent asymmetries at LHCb

Frédéric Dupertuis
Laboratoire de Physique des Hautes Energies (LPHE),
Ecole Polytechnique Fédérale de Lausanne (EPFL),
CH-1015 Lausanne, SWITZERLAND.
On behalf of the LHCb collaboration.

1 Introduction

LHCb [1] has been acquiring physics data since 2010 and recorded about 0.04 fb^{-1} in 2010 and 1.1 fb^{-1} in 2011 at center-of-mass energy ($\sqrt{s}$) of 7 TeV. In 2012, it is projected to record 2.2 fb^{-1} at an energy of $\sqrt{s} = 8$ TeV with a nominal instantaneous luminosity of $\mathcal{L} = 4 \times 10^{32}\ \mathrm{cm}^{-2}\,\mathrm{s}^{-1}$, before a long shutdown of almost two years. The data taking is expected to resume by the end of 2014 at $\sqrt{s} = 13 - 14$ TeV, before a second long shutdown in 2018 when the upgraded LHCb detector components will be installed. Since the $b\bar{b}$ cross section depends almost linearly on $\sqrt{s}$, this will lead to an increase of about 100% in $b\bar{b}$ pairs yield at $\sqrt{s} = 14$ compared to $\sqrt{s} = 7$. By 2018, a data sample larger than 8 fb^{-1} is expected to have been recorded, leading to an increase of about a factor four in statistical power with respect to the 1 fb^{-1} sample recorded at $\sqrt{s} = 7$ TeV.

The LHCb upgrade [2] is designed to take data up to a luminosity of $\mathcal{L} = 2 \cdot 10^{33}\,\mathrm{cm}^{-2}\,\mathrm{s}^{-1}$ at $\sqrt{s} = 13 - 14$ TeV, recording more than 5 fb^{-1} each year. In order not to suffer from large pile-up, the 25 ns bunch spacing of the LHC will be required. The detector readout will be upgraded to allow the full 40 MHz LHC interaction rate to be read into a software trigger, improving the trigger efficiency on hadronic modes a factor 2 (Figure 1). With an operation time of 10 years starting from 2019, it is expected to record more than 50 fb^{-1}. That will lead to a gain in statistical power by a factor ten with respect to the 1 fb^{-1} of LHCb. This paper are based on the LHCb upgrade Letter-Of-Intent (LoI) [2], the LHCb upgrade Framework TDR [3] and the LHCb prospects paper [4]. A summary of the status and the prospects of the time-dependent CP-observables is provided with their statistical error expectations. Systematic errors are expected to be kept below the statistical ones throughout the upgrade programme.

2 Weak phase in tree-dominant $b \to c\bar{c}s$ transitions

B^0/B^0_s-mesons decaying to CP-eigenstates through tree-dominant $b \to c\bar{c}s$ transitions give rise to a weak phase, coming from the interference between the direct decay and the decay after mixing. For $B^0_s \to J/\psi\phi$ and $B^0_s \to J/\psi\pi\pi$ decays the weak phase



Figure 1: *Left*: Trigger yield as function of the instantaneous luminosity for different decay channels with the current LHCb trigger system. The trigger yield for hadronic modes ($\pi\pi, \phi\gamma, D_sK$) flatens compared to modes with muons in their final states ($\psi\phi$). *Right*: Global fit of $\Delta\Gamma_s - \phi_s$ results performed by HFAG including the preliminary results of LHCb with 1 fb^{-1} of data.

ϕ_s is equal to $\phi_{s,M} - 2\,\phi_{s,D}$ where $\phi_{s,M}$, $\phi_{s,D}$ are the weak phases rising in the mixing and decay, respectively. Within the SM, $\phi_{s,M}^{\rm SM} = -2\beta_s$ and $\phi_{s,D}^{\rm SM} = \phi_s^{\rm SM,penguin}$ where β_s is the weak phase associated to the CKM element V_{ts}, which is predictaed to be small $-2\beta_s = (-0.036 \pm 0.002)$ rad [4], and $\phi_s^{\rm SM,penguin}$ is the phase coming from the doubly Cabibbo suppressed penguin decay involving V_{ub} which is expected to be even smaller than β_s. In the presence of New Physics (NP), the measured weak phase ϕ_s can be significantly enlarged with respect to the very small contribution from the SM. The full tagged angular time-dependent analysis of $B_s^0 \to J/\psi\phi$ and tagged time-dependent analysis of $B_s^0 \to J/\psi\pi\pi$ allow a precision measurement of ϕ_s. With 1 fb^{-1} of data, results of ϕ_s from a preliminary analysis of $B_s^0 \to J/\psi\phi$ decays and from $B_s^0 \to J/\psi\pi\pi$ decays give:

$$\phi_s^{J/\psi\phi} = -0.001 \pm 0.101\,({\rm stat}) \pm 0.027\,({\rm syst})\,{\rm rad}\ [5],$$
$$\phi_s^{J/\psi\pi\pi} = -0.019\,^{+0.173}_{-0.174}\,({\rm stat})\,^{+0.004}_{-0.003}\,({\rm syst})\,{\rm rad}\ [6].$$

The combined result is $\phi_s = -0.002 \pm 0.083\,({\rm stat}) \pm 0.027\,({\rm syst})\,{\rm rad}$ [5] (Figure 1). These results are compatible with the SM prediction and more data is required to pin down possible small NP effects.

The expected precisions for measuring ϕ_s are 0.025 (0.045) rad with the full LHCb dataset and 0.008 (0.014) rad with LHCb upgrade where the theoretical uncertainty of 0.003 (0.01) rad for $B_s^0 \to J/\psi\phi$ ($B_s^0 \to J/\psi\pi\pi$) will allow sensitivity to small NP phenomena. These results will be compared to measurements of ϕ_s in different decay modes such as $B_s^0 \to J/\psi\eta$, $B_s^0 \to J/\psi\eta'$ and $B_s^0 \to D_s^+D_s^-$ which have more significant penguin contributions. With 1 fb^{-1}, preliminary results yields 477 ± 23 $B_s^0 \to D_s^+D_s^-$

candidates ($\mathcal{B}(\overline{B}_s^0 \to D_s^+ D_s^-)/\mathcal{B}(\overline{B}^0 \to D^+ D_s^-) = 0.508 \pm 0.026\,(\text{stat}) \pm 0.043\,(\text{syst})$) [7] so the ϕ_s measurement in these decays will soon commence at LHCb. A clean measurement of ϕ_s can be obtained by performing a tagged time-dependent Dalitz-plot analysis of $B_s^0 \to \overline{D}^0(\to KK)KK$ penguin-free decays. With 0.62 fb^{-1}, 104 ± 29 $B_s^0 \to \overline{D}^0(\to K\pi)KK$ candidates ($\mathcal{B} = (4.7 \pm 0.9\,(\text{stat}) \pm 0.6\,(\text{syst}) \pm 0.5\,(f_s/f_d)) \cdot 10^{-5}$) [8] have been found which would yield to about 10 $B_s^0 \to \overline{D}^0(\to KK)KK$. A measurement of ϕ_s in this channel can be attempted with the LHCb upgrade.

In the B^0 sector, a tagged time-dependent analysis of $B^0 \to J/\psi K_S^0$ decays allows a measurement of the weak phase ϕ that gives within the SM almost the CKM angle β. As for ϕ_s, a small pollution $\phi^{\text{SM,penguin}}$ from penguin diagrams is expected. New NP particles entering the loop in penguin diagrams can give rise to new phase ϕ^{NP} such that the measured $\phi = 2\beta + \phi^{\text{SM,penguin}} + \phi^{\text{NP}}$. With 1 fb^{-1}, the result of $\sin(2\beta)$ from the LHCb analysis of $B^0 \to J/\psi K_S^0$ decays was presented at the CKM workshop [9], $\sin(2\beta) = 0.73 \pm 0.07\,(\text{stat}) \pm 0.04\,(\text{syst})$ [10]. By the end of 2012, LHCb will be able to provide competitive results on the single measurement of $\sin(2\beta)$ with $B^0 \to J/\psi(\to \mu^+\mu^-)K_S^0(\to \pi^+\pi^-)$ and with the full LHCb dataset, the world's best measurement is expected. With the LHCb upgrade the statistical error on $\sin(2\beta)$ is expected to go down to 0.0035. A clean measurement of $\sin(2\beta)$ can be obtained by performing a full tagged time-dependent Dalitz-plot analysis of $B_s^0 \to \overline{D}^0(\to KK)\pi\pi$ penguin-free decays. With 0.62 fb^{-1}, 8060 ± 29 $B_s^0 \to \overline{D}^0(\to K\pi)\pi\pi$ candidates [8] have been found which would translate to about 800 $B_s^0 \to \overline{D}^0(\to KK)\pi\pi$.

The penguin pollution in $b \to c\bar{c}s$ transitions is expected to be smaller than the weak phase arising from the tree processes within the SM. But with the precision that LHCb is aiming at for ϕ_s and $\sin(2\beta)$, and furthermore with the LHCb upgrade, it becomes mandatory to have a handle on the weak phase coming from the penguin process. This can be assessed by performing the analysis of SU(3)-related decays such as $B_s^0 \to J/\psi K^*$, $B^0 \to J/\psi\rho^0$ or $B^0 \to J/\psi\phi$ in the case of $B_s^0 \to J/\psi\phi$, and $B_s^0 \to J/\psi K_S^0$ or $B^0 \to J/\psi\pi^0$ channels in the case of $B^0 \to J/\psi K_S^0$. With 0.37 fb^{-1}, 116 ± 13 $B_s^0 \to J/\psi K_S^0$ candidates ($\mathcal{B} = (1.83 \pm 0.21\,(\text{stat}) \pm 0.10\,(\text{syst}) \pm 0.14\,(f_s/f_d) \pm 0.07\,(\mathcal{B}(B^0 \to J/\psi K^0))) \cdot 10^{-5}$) [11] have been observed and with 0.41 fb^{-1}, 114 ± 11 $B_s^0 \to J/\psi K^*$ candidates ($\mathcal{B} = (4.4\,^{+0.5}_{-0.4}\,(\text{stat}) \pm 0.8\,(\text{syst})) \cdot 10^{-5}$) [12] have been observed. This means that with complete analysis of these channels, LHCb will be able to provide first constraints on the penguin pollution size with the dataset recorded up to the end of 2012.

3 Weak phase in penguin-only $b \to sq\bar{q}$ with $q = s, d$ transitions

In the B_s^0 sector, $b \to sq\bar{q}$ transitions with $q = s, d$ are penguin-only processes. With the dominant t-loop there is a cancellation between the weak phase in the mixing and in the decay, so the measured weak phase ϕ_s^{eff} vanishes within the SM. ϕ_s^{eff} can be extracted from a tagged angular time-dependent analysis of $B_s^0 \to \phi\phi$ and $B_s^0 \to K^{*0}\overline{K}^{*0}$ decays and a tagged time-dependent analysis of $B_s^0 \to \phi f_0(980)$.

An untagged angular time-integrated analysis of $B_s^0 \to \phi\phi$ candidates has been performed to measure the triple product asymmetries A_U, A_V where 801 ± 29 $B_s^0 \to \phi\phi$ have been observed with a negligible S-wave fraction of $(1.3 \pm 1.2)\%$ [13]. Compatible with a vanishing ϕ_s^{eff} they have been measured with:

$$\begin{aligned} A_U &= -0.055 \pm 0.036\,(\text{stat}) \pm 0.018\,(\text{syst})\ [13], \\ A_V &= +0.010 \pm 0.036\,(\text{stat}) \pm 0.018\,(\text{syst})\ [13]. \end{aligned}$$

With 36 pb^{-1}, 50 ± 7 $B_s^0 \to K^{*0}\overline{K}^{*0}$ candidates ($\mathcal{B} = (2.81 \pm 0.46\,(\text{stat}) \pm 0.45\,(\text{syst}) \pm 0.34\,(f_s/f_d)) \cdot 10^{-5}$) have been found with a CP-averaged K^{*0} longitudinal polarisation fraction of $f_L = 0.31 \pm 0.12\,(\text{stat}) \pm 0.04\,(\text{syst})$ [14]. A tagged angular time-dependent analyses of these decays are expected with the 2011-2012 dataset and the precision on ϕ_s^{eff} with 1 fb^{-1} has been estimated to be around 0.3-0.4 rad with the channel $B_s^0 \to \phi\phi$ and 0.03 rad from 50 fb^{-1} of data with the LHCb upgrade experiment.

In the B^0 sector, the weak phase $\sin(2\beta^{\text{eff}})$ rising from $b \to sq\overline{q}$ processes with $q = s, d$ is not far from $\sin(2\beta)$ within the SM, with $|\Delta S| = |\sin(2\beta^{\text{eff}}) - \sin(2\beta)| \lesssim 0.1$ [4]. The current single best measurements of $\sin(2\beta^{\text{eff}})$ come from the B-factories with a precision of about 0.08-0.10 using $B^0 \to \eta' K_S^0$ decays. LHCb is currently studying to the $B^0 \to \phi K_S^0$ decays and the precision on $\sin(2\beta^{\text{eff}})$ is expected to go down to 0.06 with 50 fb^{-1}.

4 Weak phase $\gamma - 2\beta_s$ with $B_s^0 \to D_s^{\mp} K^{\pm}$

Performing a tagged time-dependent analysis of $B_s^0 \to D_s^{\mp} K^{\pm}$ decays allows the weak phase $\gamma - 2\beta_s$ to be measured, where $\gamma = \arg[-V_{ud}V_{ub}^*/(V_{cd}V_{cb}^*)]$. First preliminary results of a fit to the CP-observables of $B_s^0 \to D_s^{\mp} K^{\pm}$ with 1 fb^{-1} have been presented at the CKM workshop [15, 16]. The precision on γ from $B_s^0 \to D_s^{\mp} K^{\pm}$ is expected to be about 11° with the full LHCb dataset and 2° with the LHCb upgrade. $B_s^0 \to D_s^{\mp} K^{\pm} \pi^+ \pi^-$ decays are also expected to provide constraints on γ.

5 Decay width difference $\Delta\Gamma_s$ and $\Delta\Gamma_d$

The best measurement of the decay width difference $\Delta\Gamma_s$ comes from the preliminary results of the tagged angular time-dependent analysis of $B_s^0 \to J/\psi\phi$ decays with 1 fb^{-1} which gives $\Delta\Gamma_s = 0.116 \pm 0.018\,(\text{stat}) \pm 0.006\,(\text{syst})$ ps^{-1} [5]. The error on $\Delta\Gamma_s$ is expected to go down to 0.003 ps^{-1} with 50 fb^{-1}. It is possible to constrain $\Delta\Gamma_s - \phi_s$ by combining the effective lifetimes of $B_s^0 \to J/\psi f_0(980)$ (CP-even) and $B_s^0 \to K^+K^-$ (CP-odd). The sign of $\Delta\Gamma_s$ has been resolved at 4.7σ to be positive using the running of the phase difference between the P-wave and S-wave around the $\phi(1020)$ resonance [17].

LHCb expects to measure $\Delta\Gamma_d$ with a precision of 0.02 ps^{-1} with 1 fb^{-1} by comparing the effective lifetimes of $B^0 \to J/\psi K_S^0$ and $B^0 \to J/\psi K^*$, and 0.002 ps^{-1} with 50 fb^{-1}.

6 Semileptonic asymmetries $a^s_{\rm sl}$ and $a^d_{\rm sl}$

The flavour-specific asymmetry $a^s_{\rm sl}$ can be measured by computing the asymmetry between $B^0_s \to D^+_s \mu^- X$ and $B^0_s \to D^-_s \mu^+ X$ yields with $D^\pm_s \to \phi(\to K^+K^-)\pi^\pm$ (use of other D_s decays is foreseen). $a^d_{\rm sl}$ can be similarly measured using $B^0 \to D^\pm \mu^\mp X$ decays with $D^\pm \to K^\mp \pi^\pm \pi^\pm$. The first LHCb results of $a^s_{\rm sl}$ with 1 fb^{-1} gives:

$$a^s_{\rm sl} = (-0.24 \pm 0.54\,({\rm stat}) \pm 0.33\,({\rm syst}))\%\ [18].$$

With the full LHCb dataset, a statistical precision of about 0.06% is expected and 0.02% with 50 fb^{-1}.



Figure 2: $a^s_{\rm sl} - a^d_{\rm sl}$ results including the LHCb results with 1 fb^{-1} [18].

7 Conclusion

Table 1 summarises the current precision and the expectations of statistical accuracy of the CP observables with the LHCb detector and the LHCb upgrade. LHCb results are already providing world's best measurements in the B^0_s sector. The LHCb upgrade will provide high precision measurements that will stringently test the CKM paradigm and contribute significantly to the search for possible small NP effects.

Parameter	LHCb 2011 (1 fb^{-1})	LHCb 2018 (8 fb^{-1})	Upgrade (50 fb^{-1})	Theory uncertainty
$\phi_s(B^0_s \to J/\psi\phi)$ [rad]	0.10	0.025	0.008	0.003
$\phi_s(B^0_s \to J/\psi\pi\pi)$ [rad]	0.17	0.045	0.014	0.01
$\sin(2\beta)(B^0 \to J/\psi K^0_S)$ [-]	0.07	-	0.0035	0.02
$\phi^{\rm eff}_s(B^0_s \to \phi\phi)$ [rad]	*0.3-0.4 (est.)*	-	0.03	< 0.02
$\sin(2\beta^{\rm eff})(B^0 \to \phi K^0_S)$ [-]	-	-	0.06	0.02
$\gamma(B^0_s \to D^\mp_s K^\pm)$ (t) [°]	-	11°	2°	negligible
$\Delta\Gamma_s(B^0_s \to J/\psi\phi)$ [ps^{-1}]	0.018	-	0.003	0.02
$\Delta\Gamma_d(B^0 \to J/\psi K^0_S/K^*)$ [ps^{-1}]	*0.02 (est.)*	-	0.002	0.001
$a^s_{\rm sl}(B^0_s \to D^\pm_s \mu^\mp X)$ [%]	0.54	0.06	0.02	0.003

Table 1: Current and expected statistical accuracy of the relevent CP-observables measurable with time-dependent analyses with LHCb and the LHCb upgrade. The current theory uncertainties are also provided. [3, 4]

References

[1] LHCb collaboration, A. A. Alves Jr *et al.*, *The LHCb detector at the LHC*, JINST **3** (2008) S08005.

[2] LHCb collaboration, *Letter of Intent for the LHCb Upgrade*, CERN-LHCC-2011-001, LHCC-I-018.

[3] LHCb collaboration, *Framework TDR for the LHCb Upgrade*, CERN-LHCC-2012-007, LHCB-TDR-012.

[4] LHCb collaboration, A. Bharucha *et al.*, *Implications of LHCb measurements and future prospects*, arXiv:1208.3355, LHCb-PUB-2012-006. And references therein.

[5] LHCb collaboration, R. Aaij *et al.*, *Tagged time-dependent angular analysis of $B_s^0 \to J/\psi\phi$ decays with 1.03 $fb^{-}1$ at LHCb*, LHCb-CONF-2012-002.

[6] LHCb collaboration, R. Aaij *et al.*, *Measurement of ϕ_s in $\overline{B}_s \to J/\psi\pi^+\pi^-$ decays*, PAPER-2012-006

[7] LHCb collaboration, R. Aaij *et al.*, *First observations and branching fraction measurements of $\overline{B}_s^0$ to double-charm final states*, LHCb-CONF-2012-009.

[8] LHCb collaboration, R. Aaij *et al.*, *Observation of the decay $B^0 \to \overline{D}^0 K^+K^-$ and evidence of $B_s^0 \to \overline{D}^0 K^+K^-$*, Phys. Rev. Lett. **109** (2012) 131801, arXiv:1207.5991.

[9] J. Wishahi, *Measurement of Δm_d, Δm_s and* $\sin(2\beta)$, Proceedings for the 2012 7th CKM International Workshop on the CKM Unitarity Triangle.

[10] LHCb collaboration, R. Aaij *et al.*, *Measurement of the time-dependent CP asymmetry in $B^0 \to J/\psi K_S^0$ decays*, arXiv:1211.6093.

[11] LHCb collaboration, R. Aaij *et al.*, *Measurement of the $B_s^0 \to J/\psi K_s^0$ branching fraction*, arXiv:1205.0934.

[12] LHCb collaboration, R. Aaij *et al.*, *Measurement of $B_s \to J/\psi\overline{K}^{*0}$ branching fraction and angular amplitudes*, Phys. Rev. **D86** (2012) 071102, arXiv:1208.0738.

[13] LHCb collaboration, R. Aaij *et al.*, *Measurement of the polarization amplitudes and triple product asymmetries in the $B_s^0 \to \phi\phi$ decay*, Phys. Lett. **B713** (2012) 369, arXiv:1204.2813.

[14] LHCb collaboration, R. Aaij *et al.*, *First observation of the decay $B_s^0 \to K^{*0}\overline{K}^{*0}$*, Phys. Lett. **B709** (2012) 50, arXiv:1111.4183.

[15] S. Blusk, *Measurement of CKM angle gamma in $B_s^0 \to D_s^{\mp}K^{\pm}$*, Proceedings for the 2012 7th CKM International Workshop on the CKM Unitarity Triangle.

[16] LHCb collaboration, R. Aaij *et al.*, *Measurement of the time-dependent CP-violation parameters in* $B_s^0 \to D_s^{\mp} K^{\pm}$, LHCb-CONF-2012-029.

[17] LHCb collaboration, R. Aaij *et al.*, *Determination of the sign of the decay width difference in the* B_s^0 *system*, Phys. Rev. Lett. **108** (2012) 241801, arXiv:1202.4717.

[18] LHCb collaboration, R. Aaij *et al.*, *Measurement of the flavour-specific CP violating asymmetry* $a_{\rm sl}^s$ *in* B_s^0 *decays*, LHCb-CONF-2012-022.